

Manual responses to visual stimuli: early and late facilitatory effects due to the offset of a peripheral cue

Respostas manuais a estímulos visuais: efeitos facilitatórios precoces e tardios relacionados ao desaparecimento de uma dica periférica

Walter Machado-Pinheiro^{1,2}
Luiz G. Gawryszewski²
Antônio Pereira Jr.³

ABSTRACT

Manual and saccadic reaction times (MRTs and SRTs) are reduced when a warning signal precedes the onset of a target. The decreasing on SRTs observed after the offset of a fixation point has been called the gap effect. Different theories have been proposed to explain it. According to some authors, the offset also allows the saccadic system to generate a separate population of SRTs, the express saccades. Nevertheless there is no agreement about the influence of the offset of a peripheral stimulus on MRT. In two experiments we tested the effects of a peripheral visual offset used as preparatory signal on MRTs to a target after variable intervals. We found a reduction on MRT at short (200-300 ms) and long (1300-2000 ms) intervals after the peripheral offset. MRT distribution shifted toward short latencies, which sometimes formed a separate population. Since MRTs obtained at long intervals were affected by the introduction of catch trials, while MRTs at short intervals were not, we propose that two different mechanisms are involved in the decreasing of MRTs: warning and temporal expectancy. Our data support the hypothesis that the temporal component involved with the preparatory stages for motor responses can be shared by saccadic movements and key press responses, allowing the reduction on motor latencies after the visual offset in the gap paradigm. Our data corroborate the three components model for the gap effect. In our view, the question of the existence or not of a gap effect for manual responses is essentially conceptual.

Keywords: Reaction time; Preparation; Expectancy; Gap effect; Visual offset; Express responses

INTRODUCTION

Since the last century there has been general agreement that reaction times are reduced when a warning signal precedes the onset of a target (for details⁽¹⁾). Ross and Ross⁽²⁻³⁾ showed that manual reaction times (MRTs) to a visual target were equally reduced no matter if the preparatory signal (S1) was the onset or the offset of a foveal stimulus. In contrast, the reduction on saccadic reaction times (SRTs) was greater when the offset was used instead of the onset of a visual preparatory stimulus⁽³⁾. However, data obtained by our group⁽⁴⁻⁵⁾ have shown that MRTs are also differentially affected by the previous onset or offset of a visual stimulus. In one of these papers⁽⁵⁾ onset and offset of a peripheral S1 were used as a warning signal. Our results showed that S1 offset produced MRTs shorter than those observed when the onset of S1 preceded the target (S2). This facilitation on

¹ Departamento de Fisiologia e Farmacologia, Instituto Biomédico, Universidade Federal Fluminense; ² Departamento de Neurobiologia, Instituto de Biologia, Universidade Federal Fluminense; ³ Laboratório de Neuroanatomia Funcional, Departamento de Fisiologia, Universidade Federal do Pará. Correspondence concerning this article should be addressed to Luiz G. Gawryszewski, Departamento de Neurobiologia, Universidade Federal Fluminense, Caixa Postal 100.180, CEP 24001-970, Niterói, RJ, Brasil (email: gnelggw@vm.uff.br); phone 55-21-717-1161 / Fax: 55-21-2719- 5934 / 55-21-2620-3100.

MRTs for offset trials was observed for S2 presented both contra- and ipsilaterally to S1. In short, our data support the idea that the onset or offset of a visual non-informative cue produce different effects on MRT to a second stimulus.

The interest in the different effects of S1 onset or offset on motor responses to S2 has increased with the studies about phenomena such as the gap effect and the express saccades. Saslow⁽⁶⁾ was the first to show that if the fixation point (FP) disappeared prior to target onset (gap paradigm) SRTs were reduced. When the FP remained on throughout the trial (overlap paradigm), the mean SRT was about 250 ms; when the FP offset was simultaneous to the stimulus onset (gap 0), the mean SRT was about 200 ms; and when the FP disappeared 200 ms before the target onset (gap paradigm), SRTs decreased to about 150 ms. This reduction on SRTs has been called "the gap effect" (for details⁽⁷⁾).

Fischer and Boch⁽⁸⁾ in monkeys and Fischer and Ramsperger⁽⁹⁾ in humans confirmed Saslow's findings regarding the gap paradigm and showed additionally that SRT distribution in many subjects can form two distinct modes or peaks. In humans, the first peak occurred around 100-120 ms and the second one occurred around 160-180 ms. The very short latency group in the SRT distribution was called Express Saccades - ESs⁽⁷⁾. According to Fischer, "express saccades are defined by their extremely short reaction times (70 ms in monkey, 100 ms in man), which often form a separate peak in the reaction time distribution"⁽⁷⁾ (p. 553) - bimodal distribution criterion. Moreover, after training, some subjects were able to reduce or even eliminate the later peaks, thus resulting in a unimodal distribution with a single peak around 100 ms⁽⁷⁾ - absolute latency criterion. Other researchers⁽¹⁰⁻¹⁴⁾, however, though confirming the reduction on SRTs with the gap paradigm, do not support the idea of ESs. According to their interpretation, neither the bimodality nor the latency criterion are convincing enough to propose the existence of ESs.

Some important additional findings were reported by Mayfrank et al.⁽¹⁵⁾ and Braun and Breitmeyer⁽¹⁶⁾. In these studies, the onset of the target was preceded by the offset of a peripheral visual cue, while FP remained unchanged, i.e. remained on. Both studies showed that SRTs were reduced and the ESs demonstrated after a peripheral offset, even with no FP offset. In agreement with Saslow⁽⁶⁾, they also showed that the best interval to produce the reduction on SRTs was between 200 and 300 ms.

The mechanisms underlying the reduction of SRTs in the gap paradigm are not yet completely understood (for a review⁽⁷⁾). It is also not clear whether the reduction on reaction times observed with visual offset represents a general sensory-motor phenomenon or if they are restricted to the saccadic system. In fact, some authors⁽¹⁷⁻²⁰⁾ have tried to elicit express manual responses using choice reaction time paradigms. In short, no evidence has been presented for the existence of a sub-population of manual reaction times corresponding to the express responses. These results led us to study the effects of visual offsets on manual responses by changing some of the

experimental procedures used by other researchers. In the present study, we employed the simple reaction time paradigm, which, in our view, is best suited to demonstrate "reflex-like" manual responses. We also decided to use a peripheral offset instead of a FP offset (as Mayfrank et al.⁽¹⁵⁾ and Braun and Breitmeyer⁽¹⁶⁾ did) and, to avoid anticipatory errors and/or time-locked manual responses, we used variable intervals between the offset of S1 and the onset of S2. Thus, our experiments differ from most of the previous ones in three important ways: the use of a simple manual reaction times paradigm; the use of a peripheral stimulus offset instead of a fixation point offset; and the use of a variable array of intervals, since most of the studies used a constant interval or a blocked array of intervals (i.e., during a block of trials, the interval was constant - typically 200 ms).

EXPERIMENT 1

In the first experiment, we collected manual reaction times to visual targets after a peripheral visual offset in order to determine whether the findings observed by Mayfrank et al.⁽¹⁵⁾ and Braun and Breitmeyer⁽¹⁶⁾ concerning SRTs also apply to MRTs. Our main questions are: 1) Is there a reduction on simple MRT when a peripheral stimulus offset precedes the target onset, such as occurred for SRT? 2) If it occurs, what is the time course? 3) Is it possible to distinguish different populations of MRTs in the latency distribution, just as it has been found for SRTs?

MATERIAL AND METHODS

Subjects

Eight subjects (four males and four females) participated voluntarily in this study. All of them were right-handed according to the Edinburgh Inventory⁽²¹⁾, had normal or corrected vision, and their ages ranged from 20 to 35 years. They were undergraduate or graduated students at Federal Fluminense University, and had already participated in previous experiments in our laboratory, although they were naive with respect to the goals of the present experiment.

Apparatus

The experiments took place in a sound-attenuated room, under dim ambient light. The subjects sat in front of a CRT screen driven by a PC-486 microcomputer, which timed the stimuli and recorded the MRT. The head was positioned in a head-and-chin rest so that the distance between the eyes and the screen was approximately 57 cm. Before data collection, subjects were given some training and the importance of maintaining fixation was stressed. During this training, eye movements were monitored by the experimenter sitting behind the subject, using a suitably oriented mirror. The observer did not notice any shift in gaze directions before target presentation.

Moreover, in this experiment, the target was always presented around the fixation point (see below), i.e., in a foveal region.

Procedure

Each trial began with the simultaneous presentation of a central point (fixation point - FP) and a square (S1 - $0.5^\circ \times 0.5^\circ$) located 4° above it. After 700 ms, S1 went off and after a variable interval, the imperative stimulus (S2 - an identical square) flashed for 100 ms around the FP. The position of S2 close to the FP helped preventing eye movements. The intervals between S1 offset and the S2 onset were 0, 100, 200, 300, 500 and 2000 ms (Figure 1). These intervals had the same probability of occurrence and varied at random. The subjects were instructed to fixate at the FP and to use the offset of S1 as a cue for the onset of S2. They were also instructed to press a microswitch with the right index finger as soon as they detected S2. After each manual response, its latency (MRT) appeared, in milliseconds, on the screen for 1000 ms. Next trial began immediately after this presentation. Stimulus luminance was 11.5 cd/m^2 and background luminance was 0.2 cd/m^2 . All the subjects performed three sessions of 300 trials in separate days (approximately 30 minutes per session). The sessions consisted in four blocks of 75 trials interspersed with one or two minutes to rest. Fifty MRT for each interval were measured daily. The first session was considered to be training, and its results were not considered in analysis. MRTs shorter than 100 ms or longer than 700 ms were considered to be errors and generated a written message "anticipation" or "slow response", respectively, which remained on the screen for 1000 ms. These trials were repeated at the end of each session.




Figure 1 - Schematic representation of stimulus display; and temporal sequence of occurrence of the fixation point (FP), the first - preparatory - stimulus (S1) and the imperative stimulus (S2) for the first experiment

The medians of MRTs obtained in the second and third days were submitted to an analysis of variance (ANOVA) in which "day" (second and third) and "interval" (0, 100, 200, 300, 500 and 2000 ms) were used as within-subjects factors. We also performed a post-hoc analysis using Newman-Keuls method. The significance level adopted was $p < 0.05$.

RESULTS AND DISCUSSION

In this first experiment, the error-trials corresponded to 4.33% (208 errors in 4800 trials), and almost all of them were anticipations (MRTs shorter than 100 ms). These errors tended to be equally distributed among subjects and between second and third days. However, anticipations increased for the longest intervals: from 19 errors at the interval 200 ms to 88 errors at the interval of 2000 ms.

The ANOVA showed a significant effect of "interval" ($F(5,35) = 26.871$, $p < 0.001$) but not "day" ($F(1,7) = 0.123$, $p = 0.733$) on the MRTs. No interaction was observed between the two factors ($F(5,35) = 0.256$, $p = 0.932$). Figure 2 shows the variation of MRTs as a function of stimulus interval. The post-hoc analysis showed that manual responses were longer for the interval of 0 ms (248 ms) than for any other interval. MRTs for the intervals of 200, 300 and 2000 ms did not differ among them (213, 217 and 216 ms, respectively) and were shorter than the others. MRTs for the intervals of 100 and 500 ms did not differ (226 and 228 ms, respectively), but were statistically different from the others. Thus, two zones of facilitation for manual responses were found: one at 200-300 ms and another at the longest interval (2000 ms).

The time course of the first facilitation is quite similar to that observed in the gap paradigm for saccades (for a review⁽⁷⁾). Indeed, in order to define the best interval to reduce SRTs,




Figure 2 - Mean latency of manual reaction times (ordinate) as a function of the interval (abscissa) for the eight subjects. The error bars indicate ± 1 standard error of the mean

Mayfrank et al.⁽¹⁵⁾ and Braun and Breitmeyer⁽¹⁶⁾ tested the effects of different gap duration (intervals) on SRT. Both studies concluded that the optimal interval was 200-300 ms, and SRTs were not so fast for shorter (50 ms) or longer (500 ms) intervals. The intervals of 200-300 ms were also the most effective in generating a noticeable peak of express saccades. On both studies with SRTs, the interval was blocked, i.e., during a block of trials the interval was constant. In our experiment, the intervals were always randomly distributed during the test. Despite this fact, we observed that the intervals of 200-300 ms were also very efficient in reducing MRTs.

MRT distributions were made for the eight subjects with the data of the second and third days grouped. Therefore, there were 100 MRTs for each tested interval in the distributions. It was used a bin width of 10 ms, i.e., the first bin included MRTs from 120 to 129 ms, and the last from 310 to 319 ms. Figure 3 shows data from one subject for the six tested intervals. From 0 to 300 ms intervals the distributions are shifted toward shorter MRTs, while longer reaction times are more frequent at interval 500. Finally, there is a tendency toward shorter MRTs at interval 2000. Figure 4 shows the MRT distributions for the eight subjects for the intervals of 0,

200 or 300 and 2000 ms. All subjects showed shorter MRTs at intervals 200/300 and 2000 ms in relation to the 0 ms interval. Notice that distinct profiles of MRT distributions emerged for the eight subjects. For some subjects a population of MRTs can be seen at short latencies: subject #2 showed a unimodal distribution peaking at about 170-180 ms for intervals 300 and 2000. For subject #6 a peak occurred at 180-190 ms for interval 2000. Unimodal distributions were also present for subjects #1 and #8 (peaks at about the latencies of 200 and 210 ms) at intervals 200 and 2000, and for subjects #4 and #7 at interval 2000 (peaks at about 220 ms). MRT distribution for subjects #3 and #5 revealed almost no MRT shorter than 200 ms. In fact, for these subjects, the first peak of MRTs on their distribution was around the latency of 220 ms.




Figure 3 - Distribution of manual reaction times in the six tested intervals for a subject (subject 1). The number of manual responses - frequency (ordinate) was plotted as a function of manual reaction times (abscissa). It was used a bin width of 10 ms and there were 100 manual responses per condition. The number in the box on the upper-left of each graph represents the median of the manual reaction times (in ms) for that interval




Figure 4 - Distribution of manual reaction times in the intervals of 0, 200 or 300 and 2000 ms for the eight subjects. The number of manual responses - frequency (ordinate) was plotted as a function of manual reaction times (abscissa). It was used a bin width of 10 ms and there were 100 manual responses per condition

Our data also showed that MRT distributions were more scattered than those observed for saccadic responses. Several profiles of distributions emerged and tended to be more variable across subjects in relation to the studies with SRTs⁽¹⁶⁾. However, despite this variability, two subjects (#2 and #6) had short MRTs forming a peak at about the latency of 180 ms. For others (e.g., #1, #7 and #8), a population of MRTs was found peaking at about 200-210 ms. This population of short MRTs is completely absent at the interval of 0 ms for all subjects. In summary, leftward shifts on manual latencies were observed for short (200-300 ms) and long (2000 ms) intervals. The offset of a peripheral S1 is able to reduce and induce shifts toward short latencies on MRTs.

The second facilitatory zone, found at the interval of 2000 ms, however, is unexpected given that this later facilitatory temporal zone was not observed for saccadic eye movements. In fact, Mayfrank et al.⁽¹⁵⁾ reported a decrease in the population of ESs at the longest tested intervals (in their case, 600/800 ms). We believe that this later facilitation is due to a more general effect related to the expectancy, by the subject, that a stimulus which did not occur at short intervals will certainly appear later, at 2000 ms. According to this point of view, expectancy (the preparatory mechanism that takes place when someone is waiting for a stimulus that will certainly occur) would have a major influence at the longest interval, but little or no effect at short intervals. This hypothesis was tested in a second experiment. Moreover, to be sure that the effect observed at long intervals was not due to the anticipation of the exact moment of target occurrence, we will vary the longest interval from 1300 to 1900 ms.

EXPERIMENT 2

In this experiment we tested the sensibility of the two facilitatory zones observed in experiment 1 to the introduction of catch trials. As mentioned above, if expectancy has a major effect on MRTs at the longest interval, the uncertainty about S2 occurrence (introduced by catch trials) would produce a major effect at the longest interval, increasing MRTs.

MATERIAL AND METHODS

Subjects

Other eight subjects (four male and four females) participated voluntarily in this experiment. All of them were right-handed according to the Edinburgh Inventory⁽²¹⁾ and had normal or corrected vision. They were undergraduate students at Federal Fluminense University, their ages ranged from 19 to 21 years, and they had never participated in previous experiments.

Apparatus

The apparatus was the same as in experiment 1, except that this experiment ran on a Apple IIe microcomputer, the stimulus

luminance was 7.5 cd/m² and the background luminance was 0.1 cd/m².

Procedure

Each trial began with the simultaneous presentation of a central point (fixation point - FP) and a square (S1 - 0.5° x 0.5°) located 4° to the right of the central point. After 1000 ms, S1 went off and after a variable interval, the imperative stimulus (an identical square - S2) flashed for 100 ms around the central point. The interval between the offset of S1 and the onset of S2 varied randomly and with the same probability among 50, 250 and 1600 ms (Figure 5). In addition, the interval of 1600 ms varied randomly from 1300 to 1900 ms. The subjects were instructed to fixate the FP and to use the offset of S1 as a cue for the onset of S2. They were also instructed to press a microswitch with the right index finger as soon as they detected S2. Subjects were notified that there were two different conditions on separate days: no-catch and catch conditions. In the no-catch condition subjects were informed that S2 would occur in 100% of the trials. In the catch condition, S2 would occur only 80% of the time. Therefore, in the latter, subjects were never sure about S2 occurrence. After each manual response the latency of that response (MRT) in milliseconds appeared on the screen for 1500 ms. Next trial began immediately after this presentation. All the subjects performed three sessions of 300 trials in separate days (approximately 30 minutes per session). Each session had four blocks of 75 trials and one or two minutes of resting between the blocks. The first session was considered training and its data were excluded from the statistical analysis. This session included separately one block of the catch condition and another of no-catch condition. In the second and third sessions, only one condition was used at a time. For 50% of the subjects the second session was performed on the catch condition and for the other 50%, the no-catch condition was used instead. MRT




Figure 5 - Schematic representation of stimulus display; and temporal sequence of occurrence of fixation point (FP), the first - preparatory - stimulus (S1) and the imperative stimulus (S2) for the second experiment

Ross and Ross onset and offset were blocked. It means that in a block of trials only onset trials were adopted; offset trials were used in another block. In fact, in another experiment⁽²²⁾ we found some evidences that the context in which offset and onset trials were presented (blocked vs non-blocked) does have an important role in determining MRT. Differences between onset and offset trials for MRTs were greater when non-blocked procedures were adopted.

In this study, the time course of the facilitation found for MRTs was similar to those obtained by Mayfrank et al.⁽¹⁵⁾, Braun and Breitmeyer⁽¹⁶⁾ and Saslow⁽⁶⁾ for SRTs. The best interval to produce facilitation on motor responses was 200-300 ms. The similarity between our data and those obtained for saccades at short intervals suggest that both motor systems must share some common mechanisms. However, differently from the studies with SRTs, we also found a second facilitatory zone at longer intervals.

Based on their data, Kingstone and Klein⁽¹³⁾, Klein et al.⁽²³⁾ and Reuter-Lorenz et al.⁽¹¹⁾ proposed the so called two-component model to explain the gap effect. According to them, the reduction on motor responses observed in the gap paradigm would have two components: First, a warning component which follows any warning event (e.g., the offset of a visual stimuli) and is common for both saccadic and manual responses; second, a fixation offset component which is specific for saccades and is observed only when FP is turned off. This second component would be mediated by the fixation neurons (FN) located in the rostral pole of superior colliculus⁽²³⁾. According to this view, the offset of a foveal stimulus would decrease the inhibitory activity that FNs exert on collicular neurons involved with eye movements, decreasing saccades latency⁽²⁴⁻²⁸⁾. However, any involvement of collicular FNs with the observed reduction on MRTs is difficult to accept for two reasons: i) There is no evidence that the offset of a peripheral stimulus can also reduce the activity of FNs; and ii) the superior colliculus has little or nothing to do with movement of distal muscles, which were used in our procedures.

More recently, Forbes and Klein⁽²⁹⁾, Taylor et al.⁽³⁰⁾ and Trappenberg et al.⁽³¹⁾ introduced a third component to this hypothesis: an endogenous top-down interference due to some instructional or temporal prediction (previsibility) of S2 occurrence. In fact, Findlay and Walker⁽³²⁾ have also emphasized the involvement of the temporal component (the "when" component) in their model for saccade generation. We propose that the decreasing on MRTs observed in our experiments can also be attributed to this endogenous top-down effect.

In our view this endogenous top-down component is related to the preparatory mechanisms that follow the peripheral offset. In the case of SRTs experiments, preparation would be represented by the increased activity of build-up neurons of superior colliculus during the gap interval (for details about build-up activity in superior colliculus⁽³³⁻³⁶⁾). For MRTs, this motor preparation can also occur and maybe contribute to a decrease on motor latencies. In fact, using experimental procedures that involve an instructed delay period between a cue

and the triggered movement, many papers have documented the effects of this motor preparation activity. This kind of activity has been shown to occur in many motor centers: the motor cortex⁽³⁷⁻³⁹⁾, the basal ganglia⁽⁴⁰⁻⁴²⁾ and also at spinal interneurons⁽⁴³⁾. It seems obvious that for the development of this preparatory activity subjects must be able to predict the timing of target occurrence - temporal expectancy. We propose that the warning effect and the temporal expectancy effect are the major responsible for MRT decreasing observed in our experiments.

The involvement of expectancy in our results can be strongly suggested by the results of the second experiment. Only MRTs at the longest interval were affected by the introduction of catch trials. We believe that when catch trials were introduced (experiment 2), volunteers could not be sure about S2 occurrence, so their MRTs increased due to a less efficient preparatory activity.

The term expectancy has been used here as a synonym for motor preparation or preparatory set. The effects of expectancy/motor preparation in decreasing responses latencies and increasing performance are well known⁽⁴⁴⁻⁴⁹⁾. Some authors consider the Contingent Negative Variation (CNV) as an electroencephalographic (EEG) representation of these phenomena^(45,47,50-51). An important finding in this matter was obtained by Gomez et al.⁽⁵²⁾. These authors found differences in cortical potentials between gap and non-gap (gap 0) paradigms. Using FP offset, they collected choice MRTs for gap-200 ms and gap-0 ms trials and showed that gap-200 induced some alterations in event-related potential which are compatible with expectancy of target occurrence by the subject. This finding corroborates the endogenous top-down influences proposed by Forbes and Klein⁽²⁹⁾, Taylor et al.⁽³⁰⁾ and Trappenberg et al.⁽³¹⁾. It means that some cortical preparation really occur during the gap interval, and it has to do with the timing of target occurrence - temporal expectancy.

An interesting aspect of expectancy is that it could also be present at short intervals in a blocked array of intervals. When just one interval is used in a block of trials, it seems obvious that subjects will develop an expectancy to the timing of stimulus onset. We believe that in experiments that use fixed intervals, specially those without catch trials, part of the facilitation found in the gap paradigm may be due to this general effect called expectancy, since under these conditions subjects can be sure about the moment of the stimulus onset.

In the study of Braun and Breitmeyer⁽¹⁶⁾, using experimental procedures similar to those adopted for us, a population of express saccades was easily found. In the SRT distribution, they found a population of saccadic responses around the latency of 100-120 ms when the interval between the peripheral offset and the stimulus onset was 200-300 ms. They comment that for all (four) tested subjects, SRT distributions showed a similar pattern. In our study, MRT distribution showed shifts toward short latencies for intervals 200, 300 and 2000. However, data obtained in this and another study⁽⁵³⁾ showed that distribution of MRTs tended to be more scattered than

those observed for SRTs⁽¹⁷⁾. Despite this, for two subjects (#2 and #6), a peak of MRTs with short latencies (at about 180 ms) emerged. For other subjects (#1, #4, #7 and #8) a peak of MRTs appeared at about 200-210 ms. Therefore, our data indicate that MRTs with short latencies (similar to the “Express Saccades” - “express MRT”) can also be produced, at least for some subjects. As we cannot propose that fixation offset effect contributes to the reduction on MRTs observed here (see above), we are forced to conclude that warning and temporal expectancy are sufficient to induce leftward shifts in our distributions, forming a separate peak of short MRTs.

An intriguing finding of the present study is the fact that MRTs observed at short intervals were not affected by catch trials. These data apparently contrast with those obtained by Jüttner and Wolf⁽⁵⁴⁾. In their study, fixed gap intervals were used (200 ms) and they found that SRTs increased as catch trials were introduced. However, I said “apparently” because the single interval used in their experiments was 200 ms. Thus, we can interpret their results as another evidence that only the last tested interval is affected by catch trials, and the temporal context is a fundamental point to be considered in reaction times experiments. Therefore, their data corroborate our interpretation that temporal expectancy has a major influence at the longest tested interval.

Finally, if fixation offset effect really cannot be considered for manual responses and peripheral offsets, why are MRTs for short intervals so fast as those obtained for long intervals? Remember that if the reduction on MRTs were mainly due to temporal expectancy, so MRTs should be faster for longer than short intervals. There are two possibilities to explain it: i) we cannot completely discard that fixation or peripheral offsets are able to produce little or residual influences on MRTs at short intervals, just as fixation offsets reduce SRTs. If fixation offset really has nothing to do with limb movements, how can the results obtained by Bekkering et al.⁽⁵⁵⁾ be explained? These authors found a significant gap effect for pointing and choice key-pressing movements. Moreover, other theories also admit that different motor systems (for example, manual responses) can be affected by the gap effect. This is the case of the “attentional theory” proposed by Fischer and co-workers (for details⁽⁷⁾). ii) Subjects could have adopted the strategy of generating two “expectancy waves” during the experiments, an early and a late one. The first “expectancy wave”, addressed to the short intervals, for strategic reasons, would not be affected by catch trials. Top-down influences should induce an increasing in this early expectancy due to the high probability of target occurrence at short intervals.

In conclusion, our data support the hypothesis that the temporal component involved with the preparatory stages for motor responses can be shared by saccadic movements and key press responses, allowing the reduction in the latency of motor responses after the visual offset in the gap paradigm. In our view, however, the question for the existence of the gap effect for manual responses needs first to be addressed at the

conceptual level. What does the gap effect really mean, for instance? Facilitation on MRTs after peripheral offset due to warning and preparation (endogenous top-down component) can be considered a “gap effect”? Or the gap effect should be considered a synonym of fixation offset effect? It will be thus necessary first to define “gap effect” in precise terms before concluding anything about its existence concerning other motor patterns beyond saccades.

RESUMO

Os tempos de reação manuais e sacádicos (TRMs e TRSs) são reduzidos quando um sinal de aviso precede o aparecimento do alvo. O decréscimo nos TRSs observados depois do desaparecimento do ponto de fixação tem sido chamado de efeito de intervalo. Teorias diferentes foram propostas para explicá-lo. De acordo com alguns autores, o desaparecimento também permite ao sistema sacádico gerar uma população separada de TRSs, as sacádicas expressas. No entanto, não há concordância sobre a influência do desaparecimento de um estímulo periférico no TRM. Em dois experimentos, testou-se os efeitos de um desaparecimento visual periférico empregado como um sinal preparatório para os TRMs a um alvo, após intervalos variáveis. Encontrou-se uma redução no TRM para intervalos curtos (200-300 ms) e longos (1300-2000 ms) após o desaparecimento periférico. A distribuição dos TRMs deslocou-se para latências curtas, formando por vezes populações separadas. Visto que os TRMs obtidos em intervalos longos foram afetados pela introdução de sessões capciosas, enquanto que os TRMs em intervalos curtos não o foram, propõe-se que dois mecanismos diferentes estão envolvidos no decréscimo dos TRMs: alerta e expectativa temporal. Nossos dados sustentam a hipótese de que o componente temporal envolvido com os estágios preparatórios das respostas motoras podem ser compartilhados pelos movimentos sacádicos e pelas respostas de apertar botões, permitindo a redução das latências motoras após o desaparecimento visual, dentro do paradigma do intervalo. Nossos dados corroboram o modelo de três componentes do efeito de intervalo. Em nosso ponto de vista, a questão da existência ou não do efeito de intervalo para respostas manuais é essencialmente conceitual.

Descritores: Tempo de reação; Preparação; Expectativa; Efeito de intervalo; Desaparecimento visual; Respostas expressas

ACKNOWLEDGMENTS

We would like to thank Dr. Carlos Eduardo Rocha-Miranda (Academia Brasileira de Ciências), Eliane Volchan (IBCCF-UFRJ) and Dr. Ronald Ranvaud (ICB - USP) for their valuable comments on an earlier version of the manuscript. We are also grateful to Dr. Ricardo Gattass for suggesting the second experiment. Research supported by FINEP, CAPES, CNPq, PRONEX/MCT and PROPP-UFF.

REFERENCES

1. James W. *The Principles of Psychology*. Vol. 1. Authorized edition. Dover Publications Inc., New York, 1890/1950.
2. Ross LE, Ross SM. Saccade latency and warning signals: stimulus onset, offset and change as warning events. *Percept Psychophys* 1980;27:251-7.
3. Ross SM, Ross LE. Saccade latency and warning signals: effects of auditory and visual stimulus onset and offset. *Percept Psychophys* 1981;29:429-37.
4. Gawryszewski LG, Thomaz TG, Machado-Pinheiro W, Sant'Anna AN. Onset and offset of a visual cue have different effects on manual reaction time to a visual target. *Braz J Med Biol Res* 1994;27:67-73.
5. Carreiro LRR, Gawryszewski LG, Magalhães FV, Tradardi V. Onset and offset inhibitions: effects of increase and decrease of cue luminance on manual reaction time to a visual target. *Braz J Med Biol Res* 1994;27:2645-51.
6. Saslow MG. Effects of components of displacement-step stimuli upon latency of saccadic eye movements. *J Opt Soc Am* 1967;57:1024-9.
7. Fischer B, Weber H. Express saccades and visual attention. *Behav Brain Sci* 1993;16:553-610.
8. Fischer B, Boch R. Saccadic eye movements after extremely short reaction times in the monkey. *Brain Res* 1983;260:21-6.
9. Fischer B, Ramsperger E. Human express saccades: extremely short reaction times of goal directed eye movements. *Exp Brain Res* 1984;57:191-5.
10. Reuter-Lorenz PA, Hughes HC, Fendrich R. The reduction of saccade latency by prior offset of the fixation point: An analysis of the gap effect. *Percept Psychophys* 1991;49:167-75.
11. Reuter-Lorenz PA, Oonk HM, Barnes LL, Hughes HC. Effects of warning signals and fixation point offsets on the latency of pro- versus antisaccades: implications for an interpretation of the gap effect. *Exp Brain Res* 1995;103:287-93.
12. Fendrich R, Hughes HC, Reuter-Lorenz PA. Fixation-point offsets reduce the latency of saccades to acoustic targets. *Percept Psychophys* 1991;50:383-7.
13. Kingstone A, Klein RM. What are human express saccades? *Percept Psychophys* 1993a;54:260-73.
14. Kingstone A, Klein RM. Visual offsets facilitate saccadic latency: does prediengagement of visuospatial attention mediate the gap effect? *J Exp Psychol Hum Percept Perform* 1993b;19:1251-65.
15. Mayfrank L, Mobashery M, Kimming H, Fischer B. The role of fixation and visual attention in the occurrence of express saccades in man. *Eur Arch Psychiatry Neurol Sci* 1986;235:269-75.
16. Braun D, Breitmeyer BG. Relationship between directed visual attention and saccadic reaction times. *Exp Brain Res* 1988;73:546-52.
17. Rogal L, Reible G, Fischer B. Reaction times of the eye and the hand of the monkey in a visual reach task. *Neurosci Lett* 1985;58:127-32.
18. Fischer B, Rogal L. Eye-hand-coordination in man: a reaction time study. *Biol Cybern* 1986;55:253-61.
19. Fischer B. Visually guided eye and hand movements in man. *Brain Behav Evol* 1989;33:109-12.
20. Iwasaki S. Facilitation of reaction times with gap paradigm: comparison of manual and saccadic responses. *Ergonomics* 1990;33:833-50.
21. Oldfield RC. The assessment and analysis of handedness: The Edinburgh inventory. *Neuropsychologia* 1971;9:97-113.
22. Machado-Pinheiro W. *Influência de fenômenos alertantes e atencionais na gênese das respostas manuais e sacádicas*. Doctoral's Thesis, Instituto de Ciências Biomédicas, Universidade de São Paulo, São Paulo; 1999.
23. Klein R, Taylor TL, Kingstone A. Against a role for attentional disengagement in the gap effect: A friendly amendment to Tam and Stelmach (1993). *Percept Psychophys* 1995;57(4):573-7.
24. Dorris MC, Munoz DP. A neural correlate for the gap effect on saccadic reaction times in monkey. *Journal of Neurophysiol* 1995;73(6):2558-62.
25. Dorris MC, Paré M, Munoz DP. Neuronal activity in monkey superior colliculus related to the initiation of saccadic eye movements. *J Neurosci* 1997;17(21):8566-79.
26. Munoz DP, Guitton D. Control of orienting gaze shifts by tectoreticulospinal system in the head-free cat. II. Sustained discharge during motor preparation and fixation. *J Neurophysiol* 1991;66(5):1624-41.
27. Munoz DP, Wurtz RH. Role of the rostral superior colliculus in active visual fixation and execution of express saccades. *J Neurophysiol* 1992;67(4):1000-2.
28. Munoz DP, Wurtz RH. Fixation cells in monkey superior colliculus. I. Characteristics of cell discharge. *J Neurophysiol* 1993a;70(2):559-75.
29. Forbes K, Klein RM. The magnitude of the fixation offset effect with endogenously and exogenously controlled saccades. *J Cognit Neurosci* 1996;8(4):344-52.
30. Taylor TL, Kingstone A, Klein RM. The disappearance of foveal and nonfoveal stimuli: decomposing the gap effect. *Can J Exp Psychol* 1998;52(4):192-9.
31. Trappenberg TP, Dorris MC, Munoz DP, Klein RM. A model for saccade initiation based on the competitive integration of exogenous and endogenous signals in the superior colliculus. *J Cognit Neurosci* 2001;13(2):256-71.
32. Findlay JM, Walker R. A model of saccade generation based on parallel processing and competitive inhibition. *Behav Brain Sci* 1999;22:661-721.
33. Munoz DP, Wurtz RH. Saccade Related activity in the monkey superior colliculus. I. Characteristics of burst and buildup cells. *J Neurophysiol* 1995a;73(6):2313-33.
34. Munoz DP, Wurtz RH. Saccade Related activity in the monkey superior colliculus. II. Spread of activity during saccades. *J Neurophysiol* 1995b;73(6):2334-48.
35. Paré M, Munoz DP. Saccadic reaction time in the monkey: Advanced preparation of oculomotor programs is primarily responsible for express saccade occurrence. *J Neurophysiol* 1996;76(6):3666-81.
36. Basso MA. Cognitive set and oculomotor control. *Neuron* 1998;21:665-8.
37. Schmidt EM, Jost RG, Davis KK. Cortical cell discharge patterns in anticipation of a trained movement. *Brain Res* 1974;75:309-11.
38. Tanji J, Evarts EV. Anticipatory activity of motor cortex neurons in relation to direction of an intended movement. *J Neurophysiol* 1976;39(5):1062-8.
39. Kubota K, Hamada I. Preparatory activity of monkey pyramidal tract neurons related to quick movement onset during visual tracking performance. *Brain Res* 1979;168:435-9.
40. Alexander GE. Selective neuronal discharge in monkey putamen reflects intended direction of planned limb movements. *Exp Brain Res* 1987;67:623-34.
41. Alexander GE, Crutcher MD. Preparation for movement: neural representations of intended directions in three motor areas of the monkey. *J Neurophysiol* 1990;64:133-50.
42. Apicella P, Scarnati E, Ljungberg T, Schultz W. Neuronal activity in the monkey striatum related to the expectation of predictable environmental events. *J Neurophysiol* 1992;68(3):945-60.
43. Prut Y, Fetz EE. Primate spinal interneurons show pre-movement instructed delay activity. *Nature* 1999;401:590-4.
44. Bertelson P. The time course of preparation. *Q J Exp Psychol* 1967;19:272-9.
45. Birbaumer N, Elbert T, Canavan AGM, Rockstroh B. Slow potentials of the cerebral cortex and behavior. *Physiol Rev* 1990;70:1-41.
46. Boelhouwer AJW, Scheirs JGM, Blumenthal TD. The effect of response preparation upon the early component of the human blink reflex. *J Psychophysiol* 1996;9:61-9.
47. Gaillard AW, Näätänen R. Slow potential changes and choice reaction time as function of interstimulus interval. *Acta Psychol* 1973;37:173-86.
48. Posner MI, Cohen Y. Attention and the control of movements. In: Stelmach GE, Requin J, editors, *Tutorials of Motor Behavior*. Amsterdam: North Holland Publishing Co.; 1980, p.243-58.
49. Posner MI, Klein RM, Summers J, Buggie, S. On the selection of signals. *Mem Cogn* 1973;1:2-12.
50. Posner MI (1975). *Psychobiology of attention*. In: Gazzaniga M, Blakemore C, editors, *Handbook of Psychology*. New York: Academic Press. 1975, p.441-80.
51. Posner MI. *Chronometric Exploration of Mind*. Hillsdale: Erlbaum Associates, 1978.
52. Gómez C, Atienza M, López-Mendoza D, Gómez GJ, Vázquez M. Cortical potentials during gap and non-gap paradigms using manual responses in humans. *Neurosci Lett* 1995;186:107-10.
53. Machado-Pinheiro W, Gawryszewski LG, Ribeiro-do-Valle LE. Gap effect and reaction time distribution: simple vs. choice manual responses. *Braz J Med Biol Res* 1998;31:1313-8.
54. Jüttner M, Wolf W. Human express saccades: catch trials influence the probability of their occurrence. In: Schmidt R, Zambardi, editors, *Oculomotor Control and Cognitive Processes*. North Holland: Elsevier Science Pub; 1991, p.163-75.
55. Bekkering H, Pratt J, Abrams RA. The gap effect for eye and hand movement. *Percept Psychophys* 1996;58:628-35.