Hypovitaminosis A: a hidden cause of neurotrophic keratitis

Eduardo Melani Rocha Diego Rocha Gutierrez

1. Department of Ophthalmology, Faculdade de Medicina de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, SP, Brazil.

Re: Wang EY, et al.: Global Trends in Blindness and Vision Impairment Resulting from Corneal Opacity 1984-2020: a meta-analysis (Ophthalmology. 2023; 130(8): 863-871)

To the Editor:

Wang et al. (1) presented a comprehensive metaanalysis of blindness and vision impairment resulting from corneal opacity. Wang et al. stated that the "major causes of corneal opacification include trachoma, infectious keratitis, xerophthalmia, use of traditional eye medicines, and ocular trauma". Grouping these conditions revealed the amount and complexity of corneal blindness worldwide, which was warranted and identified at a crucial time. The discussion highlighted the likely role of vitamin A deficiency in corneal opacity.

Vitamin A deficiency is detrimental to corneal transparency. However, the understanding of the mechanisms of hypovitaminosis A in keratomalacia remains poor. Furthermore, it is inaccurate to assume that reduced tear flow is the sole cause of keratomalacia. A study published in 1934 by Mellanby demonstrated a probable neurotrophic origin for vitamin A deficiency-associated corneal opacity⁽²⁾. In the study, vitamin A deprivation in rats not only induced keratomalacia but also revealed trigeminal ganglion degeneration in the histological sections. These findings support the hypothesis that vitamin A deficiency induces keratomalacia via sensorial denervation of the cornea. Therefore, hypovitaminosis A is a subtype of neurotrophic keratitis. Xerophthalmia

is estimated to blind half a million children each year⁽³⁾. Considering this mechanism, hypovitaminosis A and its clinical spectrum may count as risk factors and triggers for xerophthalmia and corneal opacities.

We would like to congratulate Wang et al. for this narrative review of corneal opacities. Additionally, we would like to propose that based on previous observations of xerophthalmia causing keratomalacia and corneal blindness due to hypovitaminosis A of different origins that xerophthalmia may be a subtype of neurotrophic keratitis. This may be attributed to the fact that vitamin A deprivation causes trigeminal ganglion degeneration.

AUTHORS' CONTRIBUTIONS:

Significant to conception and design: Eduardo Melani Rocha, Diego Rocha Gutierrez. Data acquisition: Eduardo Melani Rocha, Diego Rocha Gutierrez. Data analysis and interpretation: Eduardo Melani Rocha, Diego Rocha Gutierrez. Manuscript drafting: Eduardo Melani Rocha, Diego Rocha Gutierrez. Significant intellectual content revision of the manuscript: Eduardo Melani Rocha, Diego Rocha Gutierrez. Final approval of the submitted manuscript: Eduardo Melani Rocha, Diego Rocha Gutierrez. Statistical analysis: not applicable. Obtaining funding: not applicable. Supervision of administrative, technical, or material support: Eduardo Melani Rocha. Research group leadership: Eduardo Melani Rocha.

REFERENCES

- Wang EY, Kong X, Wolle M, Gasquet N, Ssekasanvu J, Mariotti SP, et al, Global trends in blindness and vision impairment resulting from corneal opacity 1984-2020: a meta-analysis. Ophthalmology. 2023;130(8):863-71.
- 2. Mellanby E. Xerophthalmia, trigeminal degeneration and vitamin A deficiency. J Pathol. 1934;38(3): 391-407.
- Sommer A. Vitamin A deficiency, child health, and survival. Nutrition. 1997;13(5):484-5

Submitted for publication: November 28, 2023 Accepted for publication: December 12, 2023

Funding: This study received no specific financial support.

Disclosure of potential conflicts of interest: None of the authors have any potential conflicts of interest to disclose.

Corresponding author: Diego Rocha Gutierres. E-mail: drgutierrez@hcrp.usp.br; emrocha@fmrp.usp.br