



Anterior chamber metastasis from adenocarcinoma of the uterine cervix

J. William Harbour¹ , Zélia Maria Corrêa² 

1. Ocular Oncology Service, Department of Ophthalmology, and Simmons Comprehensive Cancer Center, University of Texas Southwestern, Dallas, TX, USA.

2. Ocular Oncology Service, Bascom Palmer Eye Institute and Sylvester Comprehensive Cancer Center, University of Miami, Miller School of Medicine, Miami, FL, USA.

A 32-year-old woman with multiorgan metastatic cervix adenocarcinoma reported pain and blurred vision in the right eye (OD). Her vision was counts fingers with an intraocular pressure of 30 mmHg OD. Figure 1A depicts the fleshy vascularized iris mass with layered hyphema and pseudohypopyon. The iris mass measuring $6.9 \times 6.8 \times 3.3$ mm on 50 MHz ultrasonography is shown in Figure 1B. Choroid OD and left eye were normal. She expired prematurely despite systemic chemotherapy. Cervix cancer is the second most prevalent malignancy in women rarely metastasizing to the iris^(1,2).



REFERENCES

1. Nair AG, Asnani HT, Mehta VC, Mehta SV, Pathak RS. Metastatic adenocarcinoma of the cervix presenting as a choroidal mass: A case report and review of the literature on cervical metastases to the eye. *Indian J Ophthalmol.* 2015;63(8):674-8.
2. Kurosawa A, Sawaguchi S. Iris metastasis from squamous cell carcinoma of the uterine cervix. Case report. *Arch Ophthalmol.* 1987;105(5):618.

Submitted for publication: November 19, 2024
Accepted for publication: December 11, 2024

Funding: This study received no specific financial support.

Disclosure of potential conflicts of interest: The authors declare no potential conflicts of interest.

Corresponding author: Zelia M. Correa.
E-mail: zcorrea@med.miami.edu

Informed consent was obtained from all patients included in this study.

 This content is licensed under a Creative Commons Attribution 4.0 International License.